



Trends and Risk Factors of Metabolic Syndrome among Korean Adolescents, 2007 to 2018

Jiun Chae, Moon Young Seo, Shin-Hye Kim, Mi Jung Park

Department of Pediatrics, Inje University Sanggye Paik Hospital, Inje University College of Medicine, Seoul, Korea

Background: There is a lack of recent research on the changes in risk factors for metabolic syndrome (MetS) in the Asian pediatric population. We aimed to determine the 12-year trends in the prevalence of MetS and relevant lifestyle factors such as smoking, exercise, and calorie intake among Korean adolescents.

Methods: We investigated trends in MetS and lifestyle factors among 6,308 adolescents aged 12 to 18 years from the Korea National Health and Nutrition Examination Survey, 2007 to 2018.

Results: The prevalence of MetS was stable from 2007 to 2018 (1.7% to 2.2%). There were significant increases in the prevalence of central obesity (from 8.1% to 11.2%, $P=0.012$) and hyperglycemia (from 5.3% to 10.4%, $P<0.001$) and decreases in hypo-high-density lipoprotein (HDL)-cholesterolemia (from 22.4% to 14.8%, $P<0.001$). Total calorie intake and calorie intake from fat significantly increased ($P<0.001$), whereas calorie intake from carbohydrates significantly decreased ($P<0.001$) during the study period. The proportions of tobacco smokers and regular walkers significantly decreased from 2007 to 2018. After controlling for all covariates, total calorie intake was positively correlated with waist circumference ($P<0.05$). HDL-cholesterol was negatively associated with carbohydrate consumption ($P<0.01$) and positively associated with fat consumption ($P<0.001$). Regular walking and regular strength training were associated with lower waist circumference ($P<0.05$). Smoking was associated with lower fasting glucose levels ($P<0.01$).

Conclusion: Although the prevalence rate of MetS is stable among Korean adolescents, the prevalence of central obesity and hyperglycemia has increased greatly in the recent decade. Public education on proper dietary intake and lifestyle modification is required.

Keywords: Adolescent; Diet; Exercise; Metabolic syndrome; Smoking

INTRODUCTION

Metabolic syndrome (MetS) is characterized by clustering of risk factors such as central obesity, hyperglycemia, dyslipidemia, and hypertension [1]. The onset of MetS in adolescence often leads to MetS in adulthood, resulting in cardiovascular disease and type 2 diabetes mellitus in later life [2,3]. Therefore, it is important to understand the modifiable predictors for MetS in adolescents to reduce the morbidity in adulthood.

During the last four decades, mean body mass index (BMI)

values among children and adolescents worldwide have increased [4]. Therefore, research on the trends of MetS and its related lifestyle factors is critical. Although several studies have reported the prevalence of MetS among contemporary Korean adolescents, only few studies have reported the trends of MetS in adolescents in recent years. The few existing studies reported varying trends for MetS trends in Korea and other countries [5-8]. Furthermore, no recent studies have reported trends of MetS and its components in the adolescent population. Thus, from a public health aspect, investigating changes in the trends

Corresponding authors: Mi Jung Park <https://orcid.org/0000-0002-7202-500X>
Department of Pediatrics, Inje University Sanggye Paik Hospital, Inje University College of Medicine, 1342 Dongil-ro, Nowon-gu, Seoul 01757, Korea
E-mail: pmj@paik.ac.kr

Shin-Hye Kim <https://orcid.org/0000-0003-0413-122X>
Department of Pediatrics, Inje University Sanggye Paik Hospital, Inje University College of Medicine, 1342 Dongil-ro, Nowon-gu, Seoul 01757, Korea
E-mail: s2635@paik.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

of MetS and its associated lifestyle factors among the pediatric population is critical to control cardiovascular risk factors in the future.

We examined trends in the prevalence of MetS and its components among adolescents aged 12 to 18 years in Korea, using data from the Korean National Health and Nutrition Examination Surveys (KNHANES) 2007 to 2018, and we analyzed the correlation between each risk factor.

METHODS

Study population

The data were obtained from the KNHANES, conducted between 2007 and 2018 by the Korea Centers for Disease Control and Prevention. A total of 8,087 individuals aged 12 to 18 years participated in the survey. The study population included participants whose anthropometric/blood pressure (BP) measurements and blood laboratory test reports were available. Those who had no more than 9 hours of fasting before blood sampling, those who took medications for hypertension and dyslipidemia, and those who had a medical history of hepatic, renal, or endocrine disease were excluded. In addition, adolescents were excluded if they reported physician-diagnosed diabetes; had fasting serum glucose levels ≥ 126 mg/dL, or had glycosylated hemoglobin (HbA1c) $\geq 6.5\%$. Based on the above criteria, a total of 6,308 teenagers (3,334 boys and 2,974 girls) aged 12 to 18 years were found eligible and were included in this study.

The 2007–2009, 2010–2012, and 2013–2015 KNHANES protocol was approved by an Institutional Review Board (IRB) of the Korea Centers for Disease Control and Prevention (IRB No. 2007-02CON-04-P/2008-04EXP-01-C/2009-01CON-03-2C, 2010-02CON-21-C/2011-02CON-06-C/2012-01EXP-01-2C, and 2013-07CON-03-4C/2013-12EXP-03-5C). The 2016 to 2018 KNHANES protocol was exempted from IRB approval based on the Korean bioethics law concept. Written informed consent was obtained from all the participants and their parents. The IRB of Inje University Sanggye Paik Hospital (IRB approval number: SGPAIK 2020-04-005) approved this research protocol.

Data collection and study variables

The height and weight of each participant were measured using a stadiometer and a digital weighing scale of 0.1 cm and 0.1 kg, respectively. BMI was calculated by dividing body weight

(kg) by the square root of height (m). BMI status was determined by sex- and age-specific percentiles of BMI: normal (BMI <85th percentile), overweight (BMI ≥ 85 th percentile and <95th percentile), obesity (BMI ≥ 95 th percentile), and extreme obesity (BMI ≥ 99 th percentile). Waist circumference was measured in the standing position, at the end of expiration, at the middle point of the lower margin of the 12th rib and upper margin of the iliac crest. Central obesity was defined as having a waist circumference ≥ 90 th percentile values for adolescents aged 12 to 15 years [9], ≥ 90 cm for boys aged 16 to 18 years, and ≥ 85 cm in girls aged 16 to 18 years [10]. BP was measured three times in the sitting position, and the mean values were used in the investigation. Hypertension was defined as systolic BP ≥ 130 mm Hg or diastolic BP ≥ 85 mm Hg [9].

The household income group was defined by equalized household income quartile values (total household income divided by the square root of the income of the household members). Duration and number of walking days a week were assessed using a questionnaire. Participants were divided into two categories depending on the walking duration per week: those who walked for 150 minutes a week or more and those who walked for less than 150 minutes per week. If the participant performed muscle training for more than 2 days a week, they were classified as regular strength trainers. Smoking status was classified into three groups: nonsmoker (subjects who never smoked in the past 30 days), occasional smokers (subjects who smoked less than 14 cigarettes in the 30 days prior to the survey), and frequent smokers (subjects who smoked more than 14 cigarettes in the 30 days prior to the survey). Daily calorie intake was estimated using the 24-hour dietary recall interviews by qualified dietary staff. Participants were asked to recall the food items they ingested during the past 24 hours, and the quantities of each item. Energy intakes were estimated using Korean Nutrition Society's nutrient intake assessment software (Can-Pro 2.0). The percentage of calories contributed by a specific energy source was calculated using the following equations: percentage of calories from carbohydrates = $(4 \times \text{grams of carbohydrate} \times 100) / \text{total calories}$; percentage of calories from protein = $(4 \times \text{grams of protein} \times 100) / \text{total calories}$; percentage of calories from fats = $(9 \times \text{grams of fat} \times 100) / \text{total calories}$.

After overnight fasting (more than 9 hours), blood samples were collected from the participants. An enzymatic colorimeter process, ADIVIA1650 (Siemens, Washington, DC, USA), was used to test serum glucose, triglyceride, and high-density

lipoprotein (HDL)-cholesterol levels from 2007 to February 15, 2008, and Hitachi Automatic Analyzer 7600 (Hitachi, Tokyo, Japan) was used from February 20, 2008 to 2018. Hyperglycemia and hypertriglyceridemia were defined as fasting glucose ≥ 100 mg/dL and triglycerides ≥ 150 mg/dL, respectively [9]. Hypo-HDL-cholesterolemia was determined in adolescents aged 10 to 15 years and boys aged 16 to 18 years as HDL-cholesterol < 40 mg/dL, or in girls aged 16 to 18 years as HDL-cholesterol < 50 mg/dL [9].

Definition of metabolic syndrome

In this study, we defined MetS using the International Diabetes Federation (IDF) consensus definition for children and adolescents [9]. According to this definition, MetS is diagnosed if central obesity presents with two or more of the following four criteria: (1) hypertriglyceridemia; (2) hypo-HDL-cholesterolemia; (3) hypertension; and (4) hyperglycemia.

Statistical analysis

Data analyses were conducted using SPSS version 25.0 (IBM Inc., Chicago, IL, USA) for Windows. We performed configuration of the database and complex multistage, probability sampling design analysis. We log-transformed the concentrations of triglycerides to obtain a normal distribution. The complex samples crosstabs procedure was used for categorical or ordinal variables, and the complex samples descriptive procedure was used for continuous variables. General linear models of the complex samples were used to analyze the association between lifestyle variables and MetS components. For all statistical tests, differences were considered statistically significant if P value was < 0.05 .

RESULTS

General characteristics of subjects

The general characteristics of the study sample are described in

Table 1. General characteristics of the participants ($n=6,308$)

Characteristic	Study year				P value
	2007–2009	2010–2012	2013–2015	2016–2018	
Number	1,834	1,718	1,380	1,376	
Male sex	964 (53.5)	926 (54.1)	722 (52.1)	722 (53.3)	0.785
Age, yr	15.0 \pm 0.1	15.1 \pm 0.1	15.1 \pm 0.1	15.3 \pm 0.1	0.003
12–15	1,182 (58.6)	1,081 (54.8)	824 (54.6)	817 (51.9)	0.019
16–18	652 (41.4)	637 (45.2)	556 (45.4)	559 (48.1)	
Household income					0.008
Quartile 1	228 (13.0)	198 (14.9)	151 (11.5)	145 (11.2)	
Quartile 2	407 (23.0)	421 (29.1)	359 (27.2)	305 (23.1)	
Quartile 3	583 (31.9)	518 (28.2)	467 (33.3)	446 (32.0)	
Quartile 4	589 (32.1)	560 (27.8)	394 (28.0)	479 (33.7)	
Tobacco smokers					
Non-smokers	1,720 (92.5)	1,586 (89.8)	1,296 (92.7)	1,315 (95.1)	< 0.001
Occasional smokers	42 (2.7)	64 (4.6)	29 (2.4)	24 (1.8)	
Frequent smokers	72 (4.8)	68 (5.6)	55 (4.9)	37 (3.1)	
Regular walking	1,086 (61.1)	963 (58.1)	736 (52.9)	224 (45.7)	< 0.001
Regular strength training	457 (26.0)	418 (25.0)	326 (24.2)	334 (25.4)	0.807
Total calorie intake, kcal/day	1,991.8 \pm 24.1	2,210.1 \pm 27.9	2,207.7 \pm 30.9	2,152.6 \pm 30.2	< 0.001
% Calories from CHO	64.0 \pm 0.3	62.6 \pm 0.4	60.4 \pm 0.4	59.7 \pm 0.4	< 0.001
% Calories from protein	14.1 \pm 0.1	14.5 \pm 0.1	14.1 \pm 0.1	14.7 \pm 0.1	0.001
% Calories from fat	21.8 \pm 0.3	23.0 \pm 0.3	24.1 \pm 0.3	25.3 \pm 0.3	< 0.001

Values are presented as number (%) or mean \pm standard error. The reference population is 2007 to 2009 participants. CHO, carbohydrate.

Table 1 and Fig. 1. The mean age of the participants during 2007 to 2009 was significantly lower than that during 2006 to 2018 (15.0 years vs. 15.3 years, $P=0.003$). Smoking rates were highest from 2010 to 2012, at 10.2%, whereas they were lowest during 2016 to 2018, at 4.9%, and the rates showed a significant decreasing trend over the study period ($P<0.001$). The proportion of regular walkers was 61.1% during 2007 to 2009, whereas they were lowest during 2016 to 2018, at 45.7%, and the rates showed a significant decreasing trend over the study period ($P<0.001$). The proportion of regular walkers was 61.1% during 2007 to 2009,

which gradually declined to 45.7% during 2016 to 2018 ($P<0.001$). Regarding energy intakes, significantly higher total calorie intake was noted during 2010 to 2018 than during 2007 to 2009 (2,152.6 kcal/day vs. 1,991.8 kcal/day, $P<0.001$). While the percentage of energy intake from carbohydrates showed a marked decrease from 64.0% to 59.7% ($P<0.001$), the percentage of energy intake from fats (from 21.8% to 25.3%, $P<0.004$)




Fig. 1. Trends in lifestyle factors among Korean adolescents over 12 years (2007 to 2018). ^a $P<0.01$, ^b $P<0.001$.

Table 2. Trends in metabolic syndrome components over 12 years

Variable	Study year				<i>P</i> for trend
	2007-2009	2010-2012	2013-2015	2016-2018	
Number	1,834	1,718	1,380	1,376	
Body mass index, kg/m ²	21.1±0.1	21.1±0.1	21.5±0.1	21.2±0.1	0.037
Obesity	104 (6.0)	100 (5.7)	81 (5.7)	98 (7.3)	0.174
Central obesity	146 (8.1)	148 (9.1)	141 (11.0)	157 (11.2)	0.012
Glucose, mg/dL	88.6±0.2	88.4±0.2	90.8±0.2	91.0±0.2	<0.001
Triglycerides, mg/dL	77.8±1.0	73.2±1.0	74.7±1.0	75.5±1.0	0.341
HDL-cholesterol, mg/dL	49.0±0.3	50.4±0.3	51.3±0.3	51.3±0.3	<0.001
Systolic BP, mm Hg	105.4±0.3	107.0±0.3	108.6±0.3	108.8±0.3	<0.001
Diastolic BP, mm Hg	67.0±0.3	66.7±0.3	67.1±0.3	67.4±0.3	0.239
Hyperglycemia	94 (5.3)	80 (4.2)	123 (9.0)	147 (10.4)	<0.001
Hypertriglyceridemia	175 (9.6)	130 (8.6)	103 (7.4)	120 (7.9)	0.086
Hypo-HDL-cholesterolemia	406 (22.4)	282 (17.3)	207 (15.0)	205 (14.8)	<0.001
Hypertension	50 (3.4)	50 (3.6)	65 (4.5)	54 (3.7)	0.479
Metabolic syndrome	30 (1.7)	27 (1.9)	28 (2.2)	33 (2.2)	0.289

Values are presented as mean±standard error or number (%). The reference population is 2007 to 2009 participants. Adjusted for age, sex, and household income.

HDL, high-density lipoprotein; BP, blood pressure.




Fig. 2. Trends in the prevalence of metabolic syndrome (MetS) and its components among Korean adolescents over 12 years (2007 to 2018). Hyper-TG, hypertriglyceridemia; Hypo-HDL-c, hypo-HDL-cholesterolemia. ^a $P < 0.05$, ^b $P < 0.001$.

Table 3. Mean difference in metabolic parameters according to lifestyle factors

Variable	Tobacco smokers ^a			Regular walking ^b		Regular strength training ^b	
	Non-smoker	Occasional smoker	Frequent smoker	No	Yes	No	Yes
Number	5,971	159	232	2,345	3,009	5,572	1,805
Waist circumference, cm	71.0±0.1	70.8±0.3	71.4±0.3	71.2±0.1	70.9±0.1 ^c	71.0±0.1	71.0±0.1
Glucose, mg/dL	89.9±0.1	88.8±0.6 ^d	88.3±0.6 ^d	89.9±0.2	89.7±0.2	89.9±0.1	89.5±0.2
Triglycerides, mg/dL	75.7±1.0	72.7±1.0	79.0±1.0	77.3±1.0	74.1±1.0 ^c	76.7±1.0	73.0±1.0 ^d
HDL-cholesterol, mg/dL	50.7±0.2	51.0±0.9	49.8±0.8	50.4±0.3	50.9±0.2	50.4±0.2	51.0±0.3
Systolic BP, mm Hg	107.5±0.2	107.2±0.9	106.2±0.8	107.6±0.3	107.2±0.2	107.5±0.2	106.9±0.3
Diastolic BP, mm Hg	67.0±0.2	67.3±0.8	67.2±0.7	67.1±0.2	66.9±0.2	67.0±0.2	66.8±0.3

Values are presented as mean ± standard error.

HDL, high-density lipoprotein; BP, blood pressure.

^aAdjusted for age, sex, study year group, body mass index (BMI), household income, regular walking, and total calorie intake, ^bAdjusted for age, sex, study year group, BMI, household income, and total calorie intake, ^c $P < 0.05$, ^d $P < 0.01$.

and proteins (from 14.1% to 14.7%, $P = 0.001$) showed a small but significant increase during the study period.

Trends in MetS and its components

Trends in obesity, MetS, and its components are presented in Table 2 and Fig. 2. Although there was a small increase in the prevalence of obesity (from 6.0% to 7.3%) and MetS (from 1.7% to 2.2%), there was no statistical significance ($P > 0.05$). Regarding the individual components of MetS, the prevalence of both central obesity (from 8.1% to 11.2%, $P = 0.012$) and hyperglycemia (from 5.3% to 10.4%, $P < 0.001$) demonstrated a marked increase over the study period. On the other hand, the prevalence of hypo-HDL-cholesterol significantly decreased

from 22.4% to 14.8% ($P < 0.001$). The prevalence of hypertriglyceridemia somewhat decreased from 9.6% to 7.9%, without statistical significance.

Association between metabolic parameters and lifestyle factors

Table 3 shows the mean differences in metabolic parameters according to lifestyle factors. Both frequent (88.3 mg/dL) and occasional smokers (88.8 mg/dL) showed significantly lower serum fasting glucose levels than non-smokers (89.9 mg/dL, $P < 0.01$). However, other metabolic parameters, including waist circumference and lipid profiles, were not influenced by smoking status. Regular walkers showed substantially lower

Table 4. Association of metabolic parameters with daily calorie intake

Variable	Total calorie intake ^a , kcal/day	% Calories from CHO ^b	% Calories from protein ^b	% Calories from fat ^b
Waist circumference, cm	0.214 (0.080) ^c	-0.006 (0.067)	0.202 (0.164)	-0.057 (0.085)
Glucose, mg/dL	0.093 (0.136)	0.107 (0.107)	-0.263 (0.284)	-0.159 (0.131)
Triglycerides, mg/dL	0.995 (1.009)	1.008 (1.008)	1.039 (1.021)	0.978 (1.011)
HDL-cholesterol, mg/dL	-0.064 (0.186)	-0.560 (0.176) ^d	0.502 (0.462)	0.804 (0.222) ^e
Systolic BP, mm Hg	0.108 (0.188)	0.409 (0.158) ^c	-0.426 (0.396)	-0.426 (0.189)
Diastolic BP, mm Hg	-0.126 (0.168)	0.014 (0.148)	-0.083 (0.336)	-0.150 (0.186)

Values are presented as beta (standard error). Adjusted for age, sex, study year group, body mass index, household income, smoking, and regular walking.

CHO, carbohydrate; HDL, high-density lipoprotein; BP, blood pressure.

^aCalculated as coefficients of outcome measures per increase of 1,000 kcal intake, ^bCalculated as coefficients of outcome measures per increase of 10%, ^c $P < 0.05$, ^d $P < 0.01$, ^e $P < 0.001$.

waist circumference (70.9 cm) than non-regular walkers (71.2 cm, $P < 0.05$). Both regular and regular strength trainers had significantly lower serum triglyceride levels than those who did not exercise on a regular basis ($P < 0.05$). Other metabolic parameters were not significantly different according to smoking status and exercise practice.

The results of the correlation between metabolic parameters and daily calorie intake are shown in Table 4. Waist circumference was positively associated with total calorie intake ($P < 0.05$). Other parameters, including glucose, triglycerides, HDL-cholesterol, and BP, were not significantly associated with total calorie intake. Regarding the relationship between the source of calorie intake and metabolic parameters, the percentage of calorie intake from carbohydrates was negatively correlated with HDL-cholesterol levels ($P < 0.01$), positively correlated with systolic BP ($P < 0.05$), and not with fasting glucose levels. The percentage of calorie intake from fats showed a strong positive correlation with HDL-cholesterol levels ($P < 0.001$), but was not related to other metabolic parameters, including waist circumference. The percentage of calorie intake from protein was not significantly associated with metabolic parameters.

DISCUSSION

The present study found no significant difference in the overall prevalence of MetS among Korean teenagers between 2007 and 2018. Although the prevalence of MetS was stable during the study period, an increasing trend in central obesity and hyperglycemia and a decreasing trend in hypo-HDL-cholesterol

emia were noted. We also verified the relationship between lifestyle factors and MetS parameters. Tobacco smoking was associated with lower fasting glucose levels. Regular walking was associated with lower waist circumference and lower triglycerides levels in this population. Total calorie intake was positively related to waist circumference, and the percentage of calorie intake from carbohydrates and fats was significantly correlated with HDL-cholesterol levels in opposite directions. To the best of our knowledge, this is the first epidemiological research study to show that cigarette smoking is linked to fasting plasma glucose in teenagers.

In this study, MetS prevalence defined according to by IDF Criteria was 1.7% to 2.2% among Korean adolescents, which is similar to 2.3% in China [11], and less than 3.9% to 5.5% in the United States [12,13] and 4% in Iran [14]. Epidemiological studies have shown that in recent decades, the trends in the prevalence of MetS in the United States are stable or decreasing [5,6], whereas it is increasing in Iran [15]. A few studies have demonstrated an increasing trend of MetS among Korean adolescents in the early 2000s [5]; however, since then, there has been a lack of research on trends in MetS. This study confirms that there is little difference in MetS prevalence among Korean adolescents during the past 12 years.

We found that the obesity prevalence remained constant among Korean adolescents, similar to the data reported in high-income countries [4,16]. However, central obesity seemed to be on the rise. A similar growing trend in central obesity among teenagers has been reported in many countries including China [17,18], Brazil [19], Australia [20], and Poland [21]. In contrast, no significant change has been reported in the

United States [22]. The proportion of regular walkers decreased with an increase in central obesity during the study period, and regular walkers had lower waist circumference. This finding is consistent with growing evidence that physical inactivity is a key risk factor in pediatric obesity [23,24]. We also found that waist circumference was positively correlated with total calorie intake, showing an increasing trend over the last 12 years. These findings suggest that excessive energy intake and sedentary lifestyle are responsible for the current increase in central obesity among Korean adolescents. As adolescence is a crucial time for positive lifestyle development, adolescent education on adequate calorie intake for sex and age, and physical activity is required to prevent and control central obesity.

The negative association between physical activity, including regular walking and regular strength training, and triglyceride level was confirmed in this study. Previous studies have shown a similar correlation between physical activity and triglycerides [25,26]. However, some studies have shown no significant relationship [27,28]; therefore, further research in this area is necessary. Although the underlying mechanism is unclear, it can be postulated that physical activity promotes triglyceride consumption by skeletal muscle, thereby decreasing the lipid concentration [29,30]. An exercise-induced increase in lipoprotein lipase activity, which helps triglyceride hydrolysis, also explains the decline in triglycerides due to physical activity [31].

In this study, hypo-HDL-cholesterolemia showed a decreasing trend, accompanied by an increase in fat intake and a decrease in carbohydrate intake. Previous meta-analyses have suggested that the intake of most types of fatty acids in replacement of carbohydrates increased HDL-cholesterol, while increased intake of trans-fatty acids was associated with the reduction of HDL-cholesterol [32,33]. In addition, several interventional studies have reported that low carbohydrate intake increases HDL-cholesterol by reducing the plasma apolipoprotein B (apoB)/apoA-1 ratio and insulin resistance [34,35]. Despite strong evidence for the impact of dietary fat/carbohydrate intake on lipid profiles in adults, only a few studies have been reported on the issue [6]. HDL-cholesterol improvement in our study population, despite increased total calorie intake and decreased physical activity, is partly explained by reduced carbohydrate/increased fat intake. Our finding supports an association between fat/carbohydrate intakes and HDL-cholesterol in the pediatric population.

In this study, we found that the prevalence of hyperglycemia

among Korean teenagers is on the rise. There has been a lack of accessible prevalence estimates for hyperglycemia among adolescents. Recently, two epidemiologic studies demonstrated an increasing trend in the prevalence of prediabetes by HbA1c and serum glucose assessment and raised the need for urgent interventions among United States adolescents [36,37]. The rising prediabetes trend in United States adolescents was primarily associated with an increase in obesity [36]. We believe that the increasing trend of central obesity could mainly explain the rise in hyperglycemia in Korean teenagers. The increase in teenagers' hyperglycemia can also be explained by the decrease in physical activity and the increase in fat intakes. Physical activity increases the number and function of glucose transporter type 4 (GLUT4) in muscle and increases blood flow to the muscle, which improves insulin sensitivity [38]. Previous studies of adults have shown that high-fat diets reduce insulin sensitivity without increasing internal fat [39]. Other previous studies of adults found that a short-term high-fat diet and physical inactivity reduce insulin sensitivity, and physical activity was found to have a protective effect on the decrease of insulin sensitivity by a high-fat diet [40]. Therefore, we think that the decline in physical activity added to the increase in dietary fat might have contributed to the rise in hyperglycemia by worsening insulin resistance among Korean adolescents.

In addition, we found a significant association between tobacco smoking and higher fasting glucose levels in this study population. The relationship between blood glucose and smoking is controversial. Some studies that reported a positive relationship between smoking and serum glucose suggested that smoking might cause insulin resistance [41,42] or pancreatic β -cell impairment [43,44]. Other studies have suggested that smoking reduces serum glucose because nicotine blocks the release of inflammatory cytokines, which inhibits inflammation associated with insulin resistance and reduces hyperglycemia [45]. Additionally, smoking-induced catecholamines increase the metabolism of soluble nutrients including glucose [45]. Regardless of the controversial association between smoking and glucose, smoking remains a major risk factor for cardiovascular disease and diabetes-related complications [46,47]. Therefore, adolescent awareness on the importance of avoiding smoking should be emphasized, regardless of their blood glucose levels.

This study has several limitations. First, as this is not a longitudinal study, a causal relationship cannot be established. Sec-

ond, because data on the lifestyle factors were collected using self-reported questionnaires, biases may affect the results. Finally, the pubertal status of the participants, which may affect insulin resistance and lipid profile, was not validated in this study. Nonetheless, this is the first epidemiological study that investigated both the trends of MetS in Korean teenagers over the last 12 years and the related lifestyle factors using a nationally representative sample. We also discussed the potential mechanisms by which lifestyle factors, including smoking and sources of energy consumption, can influence the metabolic components.

In conclusion, while the prevalence of MetS has been constant over the past decade, the prevalence of central obesity and hyperglycemia has increased among Korean adolescents. Korean teenagers need public education regarding proper lifestyle modification and adequate calorie consumption.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

AUTHOR CONTRIBUTIONS

Conception or design: J.C., S.H.K., M.J.P.

Acquisition, analysis, or interpretation of data: J.C., S.H.K.

Drafting of the work or revision: J.C., M.Y.S., S.H.K., M.J.P.

Final approval of the manuscript: J.C., M.Y.S., S.H.K., M.J.P.

ORCID

Jiun Chae <https://orcid.org/0000-0003-4595-0635>

Mi Jung Park <https://orcid.org/0000-0002-7202-500X>

Shin-Hye Kim <https://orcid.org/0000-0003-0413-122X>

FUNDING

None

ACKNOWLEDGMENTS

We thank the Korea Centers for Disease Control and Prevention for providing the data.

REFERENCES

1. Kassi E, Pervanidou P, Kaltsas G, Chrousos G. Metabolic syndrome: definitions and controversies. *BMC Med* 2011;9:48.
2. Morrison JA, Friedman LA, Gray-McGuire C. Metabolic syndrome in childhood predicts adult cardiovascular disease 25 years later: the Princeton Lipid Research Clinics Follow-up Study. *Pediatrics* 2007;120:340-5.
3. Morrison JA, Friedman LA, Wang P, Glueck CJ. Metabolic syndrome in childhood predicts adult metabolic syndrome and type 2 diabetes mellitus 25 to 30 years later. *J Pediatr* 2008;152: 201-6.
4. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;390:2627-42.
5. Lim S, Jang HC, Park KS, Cho SI, Lee MG, Joung H, et al. Changes in metabolic syndrome in American and Korean youth, 1997-2008. *Pediatrics* 2013;131:e214-22.
6. Lee AM, Gurka MJ, DeBoer MD. Trends in metabolic syndrome severity and lifestyle factors among adolescents. *Pediatrics* 2016;137:e20153177.
7. Barzin M, Hosseinpanah F, Saber H, Sarbakhsh P, Nakhoda K, Azizi F. Gender differences time trends for metabolic syndrome and its components among Tehranian children and adolescents. *Cholesterol* 2012;2012:804643.
8. Park MJ, Boston BA, Oh M, Jee SH. Prevalence and trends of metabolic syndrome among Korean adolescents: from the Korean NHANES survey, 1998-2005. *J Pediatr* 2009;155:529-34.
9. International Diabetes Federation: The IDF consensus definition of the metabolic syndrome in children and adolescents. Available from: <https://www.idf.org/e-library/consensus-statements/61-idf-consensus-definition-of-metabolic-syndrome-in-children-and-adolescents.html> (cited 2021 May 7).
10. Lee SY, Park HS, Kim DJ, Han JH, Kim SM, Cho GJ, et al. Appropriate waist circumference cutoff points for central obesity in Korean adults. *Diabetes Res Clin Pract* 2007;75:72-80.
11. Zhu Y, Zheng H, Zou Z, Jing J, Ma Y, Wang H, et al. Metabolic syndrome and related factors in Chinese children and adolescents: analysis from a Chinese national study. *J Atheroscler Thromb* 2020;27:534-44.
12. Ford ES, Li C, Zhao G, Pearson WS, Mokdad AH. Prevalence of the metabolic syndrome among U.S. adolescents using the definition from the International Diabetes Federation. *Diabe-*

- tes Care 2008;31:587-9.
13. Park J, Hilmers DC, Mendoza JA, Stuff JE, Liu Y, Nicklas TA. Prevalence of metabolic syndrome and obesity in adolescents aged 12 to 19 years: comparison between the United States and Korea. *J Korean Med Sci* 2010;25:75-82.
 14. Kelishadi R, Heshmat R, Farzadfar F, Esmaeil Motlag M, Bahreynian M, Safiri S, et al. Prevalence of cardio-metabolic risk factors in a nationally representative sample of Iranian adolescents: the CASPIAN-III Study. *J Cardiovasc Thorac Res* 2017;9:12-20.
 15. Kelishadi R, Hovsepian S, Djalalinia S, Jamshidi F, Qorbani M. A systematic review on the prevalence of metabolic syndrome in Iranian children and adolescents. *J Res Med Sci* 2016;21:90.
 16. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:766-81.
 17. Liang YJ, Xi B, Song AQ, Liu JX, Mi J. Trends in general and abdominal obesity among Chinese children and adolescents 1993-2009. *Pediatr Obes* 2012;7:355-64.
 18. Zhang YX, Wang SR, Chen M, Cheng Y. Recent trends in body mass index and waist circumference among children and adolescents in Shandong China. *J Trop Pediatr* 2017;63:461-7.
 19. Casagrande D, Waib PH, Sgarbi JA. Increase in the prevalence of abdominal obesity in Brazilian school children (2000-2015). *Int J Pediatr Adolesc Med* 2017;4:133-7.
 20. Hardy LL, Mhrshahi S, Gale J, Drayton BA, Bauman A, Mitchell J. 30-Year trends in overweight, obesity and waist-to-height ratio by socioeconomic status in Australian children, 1985 to 2015. *Int J Obes (Lond)* 2017;41:76-82.
 21. Suder A, Gomula A, Koziel S. Central overweight and obesity in Polish schoolchildren aged 7-18 years: secular changes of waist circumference between 1966 and 2012. *Eur J Pediatr* 2017;176:909-16.
 22. Xi B, Mi J, Zhao M, Zhang T, Jia C, Li J, et al. Trends in abdominal obesity among U.S. children and adolescents. *Pediatrics* 2014;134:e334-9.
 23. Grigorakis DA, Georgoulis M, Psarra G, Tambalis KD, Panagiotakos DB, Sidossis LS. Prevalence and lifestyle determinants of central obesity in children. *Eur J Nutr* 2016;55:1923-31.
 24. El-Kassas G, Ziade F. Exploration of the risk factors of generalized and central obesity among adolescents in North Lebanon. *J Environ Public Health* 2017;2017:2879075.
 25. Jenkins GP, Evenson KR, Herring AH, Hales D, Stevens J. Cardiometabolic correlates of physical activity and sedentary patterns in U.S. youth. *Med Sci Sports Exerc* 2017;49:1826-33.
 26. LeBlanc AG, Janssen I. Dose-response relationship between physical activity and dyslipidemia in youth. *Can J Cardiol* 2010;26:201-5.
 27. Whitaker KM, Pettee Gabriel K, Buman MP, Pereira MA, Jacobs DR Jr, Reis JP, et al. Associations of accelerometer-measured sedentary time and physical activity with prospectively assessed cardiometabolic risk factors: the CARDIA study. *J Am Heart Assoc* 2019;8:e010212.
 28. Heshmat R, Qorbani M, Shahr Babaki AE, Djalalinia S, Ataei-Jafari A, Motlagh ME, et al. Joint association of screen time and physical activity with cardiometabolic risk factors in a national sample of Iranian adolescents: the CASPIANIII Study. *PLoS One* 2016;11:e0154502.
 29. Baran J, Weres A, Czenczek-Lewandowska E, Wyszynska J, Luszczki E, Deren K, et al. Blood lipid profile and body composition in a pediatric population with different levels of physical activity. *Lipids Health Dis* 2018;17:171.
 30. Mann S, Beedie C, Jimenez A. Differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations. *Sports Med* 2014;44:211-21.
 31. Wang Y, Xu D. Effects of aerobic exercise on lipids and lipoproteins. *Lipids Health Dis* 2017;16:132.
 32. Yanai H, Katsuyama H, Hamasaki H, Abe S, Tada N, Sako A. Effects of dietary fat intake on HDL metabolism. *J Clin Med Res* 2015;7:145-9.
 33. Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *Am J Clin Nutr* 2003;77:1146-55.
 34. Morton AM, Furtado JD, Mendivil CO, Sacks FM. Dietary unsaturated fat increases HDL metabolic pathways involving apoE favorable to reverse cholesterol transport. *JCI Insight* 2019;4:e124620.
 35. Desroches S, Paradis ME, Perusse M, Archer WR, Bergeron J, Couture P, et al. Apolipoprotein A-I, A-II, and VLDL-B-100 metabolism in men: comparison of a low-fat diet and a high-monounsaturated fatty acid diet. *J Lipid Res* 2004;45:2331-8.
 36. Lee AM, Fermin CR, Filipp SL, Gurka MJ, DeBoer MD. Examining trends in prediabetes and its relationship with the metabolic syndrome in US adolescents, 1999-2014. *Acta Diabetol* 2017;54:373-81.

37. Umapathi KK, Thavamani A, Al-Kindi S. Prediabetes in children and adolescents in the United States: prevalence estimates and comorbidities: a population analysis. *J Pediatr Endocrinol Metab* 2019;32:187-9.
38. Strasser B. Physical activity in obesity and metabolic syndrome. *Ann N Y Acad Sci* 2013;1281:141-59.
39. von Frankenberg AD, Marina A, Song X, Callahan HS, Kratz M, Utzschneider KM. A high-fat, high-saturated fat diet decreases insulin sensitivity without changing intra-abdominal fat in weight-stable overweight and obese adults. *Eur J Nutr* 2017;56:431-43.
40. Stettler R, Ith M, Acheson KJ, Decombaz J, Boesch C, Tappy L, et al. Interaction between dietary lipids and physical inactivity on insulin sensitivity and on intramyocellular lipids in healthy men. *Diabetes Care* 2005;28:1404-9.
41. Morimoto A, Tatsumi Y, Deura K, Mizuno S, Ohno Y, Watanabe S. Impact of cigarette smoking on impaired insulin secretion and insulin resistance in Japanese men: the Saku Study. *J Diabetes Investig* 2013;4:274-80.
42. Piatti P, Setola E, Galluccio E, Costa S, Fontana B, Stuccillo M, et al. Smoking is associated with impaired glucose regulation and a decrease in insulin sensitivity and the disposition index in first-degree relatives of type 2 diabetes subjects independently of the presence of metabolic syndrome. *Acta Diabetol* 2014;51:793-9.
43. Xu M, Zhou Y, Xu B, Sun J, Wang T, Lu J, et al. Associations of smoking and alcohol consumption with impaired β -cell function in Chinese men. *J Diabetes* 2016;8:434-41.
44. Ostgren CJ, Lindblad U, Ranstam J, Melander A, Rastam L; Skaraborg Hypertension and Diabetes Project. Associations between smoking and beta-cell function in a non-hypertensive and non-diabetic population. Skaraborg Hypertension and Diabetes Project. *Diabet Med* 2000;17:445-50.
45. Haskins AE, Bertone-Johnson ER, Pekow P, Carbone E, Fortner RT, Chasan-Taber L. Smoking during pregnancy and risk of abnormal glucose tolerance: a prospective cohort study. *BMC Pregnancy Childbirth* 2010;10:55.
46. Fa-Binefa M, Clara A, Perez-Fernandez S, Grau M, Degano IR, Marti-Lluch R, et al. Early smoking-onset age and risk of cardiovascular disease and mortality. *Prev Med* 2019;124:17-22.
47. Kelishadi R, Noori A, Qorbani M, Rahimzadeh S, Djalalinia S, Shafiee G, et al. Are active and passive smoking associated with cardiometabolic risk factors in adolescents? The CASPIAN-III Study. *Paediatr Int Child Health* 2016;36:181-8.