

SUPPLEMENTARY REFERENCES

1. Annibali G, Lucertini F, Agostini D, Vallorani L, Gioacchini A, Barbieri E, et al. Concurrent aerobic and resistance training has anti-inflammatory effects and increases both plasma and leukocyte levels of IGF-1 in late middle-aged type 2 diabetic patients. *Oxid Med Cell Longev*. 2017;2017:3937842.
2. Archer T, Fredriksson A, Schutz E, Kostrzewa RM. Influence of physical exercise on neuroimmunological functioning and health: aging and stress. *Neurotox Res* 2011;20:69-83.
3. Balducci S, Zanuso S, Cardelli P, Salerno G, Fallucca S, Nicolucci A, et al. Supervised exercise training counterbalances the adverse effects of insulin therapy in overweight/obese subjects with type 2 diabetes. *Diabetes Care* 2012;35:39-41.
4. Balducci S, Zanuso S, Cardelli P, Salvi L, Mazzitelli G, Bazuro A, et al. Changes in physical fitness predict improvements in modifiable cardiovascular risk factors independently of body weight loss in subjects with type 2 diabetes participating in the Italian Diabetes and Exercise Study (IDES). *Diabetes Care* 2012;35:1347-54.
5. Banitalebi E, Kazemi A, Faramarzi M, Nasiri S, Haghighi MM. Effects of sprint interval or combined aerobic and resistance training on myokines in overweight women with type 2 diabetes: A randomized controlled trial. *Life Sci* 2019;217:101-9.
6. Bouchi R, Sonoda N, Itoh J, Ono Y, Fukuda T, Takeuchi T, et al. Effects of intensive exercise combined with dapagliflozin on body composition in patients with type 2 diabetes: a randomized controlled trial. *Endocr J* 2021;68:329-43.
7. Brozic AP, Marzolini S, Goodman JM. Effects of an adapted cardiac rehabilitation programme on arterial stiffness in patients with type 2 diabetes without cardiac disease diagnosis. *Diab Vasc Dis Res* 2017;14:104-12.
8. Buch A, Eldor R, Kis O, Keinan-Boker L, Dunsky A, Rubin A, et al. The effect of circuit resistance training, empagliflozin or “vegeterranean diet” on physical and metabolic function in older subjects with type 2 diabetes: a study protocol for a randomized control trial (CEV-65 trial). *BMC Geriatr* 2019;19:228.
9. Chen X, Sun X, Wang C, He H. Effects of exercise on inflammatory cytokines in patients with type 2 diabetes: a meta-analysis of randomized controlled trials. *Oxid Med Cell Longev* 2020;2020:6660557.
10. Church TS, Blair SN, Cocreham S, Johannsen N, Johnson W, Kramer K, et al. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2010;304:2253-62.
11. Cobbold C. Type 2 diabetes mellitus risk and exercise: is resistin involved? *J Sports Med Phys Fitness* 2019;59:290-7.
12. Cronin O, Keohane DM, Molloy MG, Shanahan F. The effect of exercise interventions on inflammatory biomarkers in healthy, physically inactive subjects: a systematic review. *QJM* 2017;110:629-37.
13. Daly RM, Miller EG, Dunstan DW, Kerr DA, Solah V, Menzies D, et al. The effects of progressive resistance training combined with a whey-protein drink and vitamin D supplementation on glycaemic control, body composition and cardiometabolic risk factors in older adults with type 2 diabetes: study protocol for a randomized controlled trial. *Trials* 2014;15:431.
14. Donges CE, Duffield R, Drinkwater EJ. Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Med Sci Sports Exerc* 2010;42:304-13.
15. Egger A, Niederseer D, Diem G, Finkenzeller T, Ledl-Kurkowsky E, Forstner R, et al. Different types of resistance training in type 2 diabetes mellitus: effects on glycaemic control, muscle mass and strength. *Eur J Prev Cardiol* 2013;20:1051-60.
16. Calle MC, Fernandez ML. Effects of resistance training on the inflammatory response. *Nutr Res Pract* 2010;4:259-69.
17. Gordon BA, Fraser SF, Bird SR, Benson AC. Insulin sensitivity not modulated 24 to 78 h after acute resistance exercise in type 2 diabetes patients. *Diabetes Obes Metab* 2013;15:478-80.
18. Grote S, Almstedt HC, Tarleton HP. Cardiometabolic health among cancer survivors: a 13-week pilot study of a combined aerobic and resistance training program. *Oncol Nurs Forum* 2016;43:306-15.
19. Hopps E, Canino B, Caimi G. Effects of exercise on inflammation markers in type 2 diabetic subjects. *Acta Diabetol* 2011;48:183-9.
20. Johansen MY, Karstoft K, MacDonald CS, Hansen KB, Ellingsgaard H, Hartmann B, et al. Effects of an intensive lifestyle intervention on the underlying mechanisms of improved glycaemic control in individuals with type 2 diabetes: a secondary analysis of a randomised clinical trial. *Diabetologia* 2020;63:2410-22.
21. Kelly LA, Loza A, Lin X, Schroeder ET, Hughes A, Kirk A, et al. The effect of a home-based strength training program on type 2 diabetes risk in obese Latino boys. *J Pediatr Endocrinol Metab* 2015;28:315-22.
22. Kim JY, Jeon JY. Role of exercise on insulin sensitivity and beta-cell function: is exercise sufficient for the prevention of youth-onset type 2 diabetes? *Ann Pediatr Endocrinol Metab* 2020;25:

- 208-16.
23. Kohut ML, McCann DA, Russell DW, Konopka DN, Cunnick JE, Franke WD, et al. Aerobic exercise, but not flexibility/resistance exercise, reduces serum IL-18, CRP, and IL-6 independent of beta-blockers, BMI, and psychosocial factors in older adults. *Brain Behav Immun* 2006;20:201-9.
 24. Kruger K, Mooren FC, Eder K, Ringseis R. Immune and inflammatory signaling pathways in exercise and obesity. *Am J Lifestyle Med* 2014;10:268-79.
 25. Leehey DJ, Collins E, Kramer HJ, Cooper C, Butler J, McBurney C, et al. Structured exercise in obese diabetic patients with chronic kidney disease: a randomized controlled trial. *Am J Nephrol* 2016;44:54-62.
 26. Liu Y, Liu SX, Cai Y, Xie KL, Zhang WL, Zheng F. Effects of combined aerobic and resistance training on the glycolipid metabolism and inflammation levels in type 2 diabetes mellitus. *J Phys Ther Sci* 2015;27:2365-71.
 27. Loprinzi PD, Moore D, Loenneke JP. Does aerobic and resistance exercise influence episodic memory through unique mechanisms? *Brain Sci* 2020;10:913.
 28. Lucotti P, Monti LD, Setola E, Galluccio E, Gatti R, Bosi E, et al. Aerobic and resistance training effects compared to aerobic training alone in obese type 2 diabetic patients on diet treatment. *Diabetes Res Clin Pract* 2011;94:395-403.
 29. Magalhaes JP, Santos DA, Correia IR, Hetherington-Rauth M, Ribeiro R, Raposo JF, et al. Impact of combined training with different exercise intensities on inflammatory and lipid markers in type 2 diabetes: a secondary analysis from a 1-year randomized controlled trial. *Cardiovasc Diabetol* 2020;19:169.
 30. Martins RA, Neves AP, Coelho-Silva MJ, Verissimo MT, Teixeira AM. The effect of aerobic versus strength-based training on high-sensitivity C-reactive protein in older adults. *Eur J Appl Physiol* 2010;110:161-9.
 31. Melo LC, Dativo-Medeiros J, Menezes-Silva CE, Barbosa FT, de Sousa-Rodrigues CF, Rabelo LA. Physical exercise on inflammatory markers in type 2 diabetes patients: a systematic review of randomized controlled trials. *Oxid Med Cell Longev* 2017;2017:8523728.
 32. Misra A, Alappan NK, Vikram NK, Goel K, Gupta N, Mittal K, et al. Effect of supervised progressive resistance-exercise training protocol on insulin sensitivity, glycemia, lipids, and body composition in Asian Indians with type 2 diabetes. *Diabetes Care* 2008;31:1282-7.
 33. Morais Junior GS, Souza VC, Machado-Silva W, Henriques AD, Avelar GG, Perez DIV, et al. Resistance training modulates the humoral inflammatory (but not the DNA Methylation) profile of diabetic older adults using metformin. *Neuroimmunomodulation* 2019;26:208-16.
 34. Nikseresht M, Agha-Alinejad H, Azarbayjani MA, Ebrahim K. Effects of nonlinear resistance and aerobic interval training on cytokines and insulin resistance in sedentary men who are obese. *J Strength Cond Res* 2014;28:2560-8.
 35. Nimmo MA, Leggate M, Viana JL, King JA. The effect of physical activity on mediators of inflammation. *Diabetes Obes Metab* 2013;15 Suppl 3:51-60.
 36. Ploeger HE, Takken T, de Greef MH, Timmons BW. The effects of acute and chronic exercise on inflammatory markers in children and adults with a chronic inflammatory disease: a systematic review. *Exerc Immunol Rev* 2009;15:6-41.
 37. Ranasinghe C, Hills AP, Constantine GR, Finlayson G, Katalanda P, King NA. Study protocol: a randomised controlled trial of supervised resistance training versus aerobic training in Sri Lankan adults with type 2 diabetes mellitus: SL-DART study. *BMC Public Health* 2018;18:176.
 38. Hu D, Russell RD, Remash D, Greenaway T, Rattigan S, Squibb KA, et al. Are the metabolic benefits of resistance training in type 2 diabetes linked to improvements in adipose tissue microvascular blood flow? *Am J Physiol Endocrinol Metab* 2018;315:E1242-50.
 39. Ryrso CK, Thaning P, Siebenmann C, Lundby C, Lange P, Pedersen BK, et al. Effect of endurance versus resistance training on local muscle and systemic inflammation and oxidative stress in COPD. *Scand J Med Sci Sports* 2018;28:2339-48.
 40. Stefanov T, Bluher M, Vekova A, Bonova I, Tzvetkov S, Kurkschiev D, et al. Circulating chemerin decreases in response to a combined strength and endurance training. *Endocrine* 2014;45:382-91.
 41. Swift DL, Johannsen NM, Myers VH, Earnest CP, Smits JA, Blair SN, et al. The effect of exercise training modality on serum brain derived neurotrophic factor levels in individuals with type 2 diabetes. *PLoS One*. 2012;7:e42785.
 42. Thent ZC, Das S, Henry LJ. Role of exercise in the management of diabetes mellitus: the global scenario. *PLoS One* 2013;8:e80436.
 43. Touvra AM, Volaklis KA, Spassis AT, Zois CE, Douda HD, Kotsa K, et al. Combined strength and aerobic training increases transforming growth factor- β 1 in patients with type 2 diabetes. *Hormones (Athens)* 2011;10:125-30.
 44. Wasserfurth P, Nebl J, Schuchardt JP, Muller M, BoBlau TK, Kruger K, et al. Effects of exercise combined with a healthy diet

- or calanus finmarchicus oil supplementation on body composition and metabolic markers: a pilot study. *Nutrients* 2020;12:2139.
45. Gordon PL, Vannier E, Hamada K, Layne J, Hurley BF, Roubenoff R, et al. Resistance training alters cytokine gene expression in skeletal muscle of adults with type 2 diabetes. *Int J Immunopathol Pharmacol* 2006;19:739-49.
 46. Brooks N, Layne JE, Gordon PL, Roubenoff R, Nelson ME, Castaneda-Sceppa C. Strength training improves muscle quality and insulin sensitivity in Hispanic older adults with type 2 diabetes. *Int J Med Sci* 2006;4:19-27.
 47. Jorge ML, de Oliveira VN, Resende NM, Paraiso LF, Calixto A, Diniz AL, et al. The effects of aerobic, resistance, and combined exercise on metabolic control, inflammatory markers, adipocytokines, and muscle insulin signaling in patients with type 2 diabetes mellitus. *Metabolism* 2011;60:1244-52.
 48. Kadoglou NP, Fotiadis G, Athanasiadou Z, Vitta I, Lampropoulos S, Vrabas IS. The effects of resistance training on ApoB/ApoA-I ratio, Lp(a) and inflammatory markers in patients with type 2 diabetes. *Endocrine* 2012;42:561-9.
 49. Swift DL, Johannsen NM, Earnest CP, Blair SN, Church TS. Effect of exercise training modality on C-reactive protein in type 2 diabetes. *Med Sci Sports Exerc* 2012;44:1028-34.
 50. Kadoglou NP, Fotiadis G, Kapelouzou A, Kostakis A, Liapis CD, Vrabas IS. The differential anti-inflammatory effects of exercise modalities and their association with early carotid atherosclerosis progression in patients with type 2 diabetes. *Diabet Med* 2013;30:e41-50.
 51. Mavros Y, Kay S, Simpson KA, Baker MK, Wang Y, Zhao RR, et al. Reductions in C-reactive protein in older adults with type 2 diabetes are related to improvements in body composition following a randomized controlled trial of resistance training. *J Cachexia Sarcopenia Muscle* 2014;5:111-20.
 52. Hsieh PL, Tseng CH, Tseng YJ, Yang WS. Resistance training improves muscle function and cardiometabolic risks but not quality of life in older people with type 2 diabetes mellitus: a randomized controlled trial. *J Geriatr Phys Ther* 2018;41:65-76.
 53. Miller EG, Sethi P, Nowson CA, Dunstan DW, Daly RM. Effects of progressive resistance training and weight loss versus weight loss alone on inflammatory and endothelial biomarkers in older adults with type 2 diabetes. *Eur J Appl Physiol* 2017;117:1669-78.
 54. Dadrass A, Mohamadzadeh Salamat K, Hamidi K, Azizbeigi K. Anti-inflammatory effects of vitamin D and resistance training in men with type 2 diabetes mellitus and vitamin D deficiency: a randomized, double-blinded, placebo-controlled clinical trial. *J Diabetes Metab Disord* 2019;18:323-31.
 55. Rech A, Botton CE, Lopez P, Quincozes-Santos A, Umpierre D, Pinto RS. Effects of short-term resistance training on endothelial function and inflammation markers in elderly patients with type 2 diabetes: a randomized controlled trial. *Exp Gerontol* 2019;118:19-25.
 56. Ranasinghe C, Devage S, Constantine GR, Katulanda P, Hills AP, King NA. Glycemic and cardiometabolic effects of exercise in South Asian Sri Lankans with type 2 diabetes mellitus: a randomized controlled trial Sri Lanka diabetes aerobic and resistance training study (SL-DARTS). *Diabetes Metab Syndr* 2021;15:77-85.
 57. Sabouri M, Hatami E, Pournemati P, Shabkhiz F. Inflammatory, antioxidant and glycemic status to different mode of high-intensity training in type 2 diabetes mellitus. *Mol Biol Rep* 2021;48:5291-304.