

**Supplementary Table 1.** Primer sequences of genes used to perform quantitative reverse transcription polymerase chain reaction

| Gene          | Forward                | Reverse               |
|---------------|------------------------|-----------------------|
| <i>18s</i>    | TCGGTCTTATTTTGCTGGTTT  | ATCGCTTGGCATCGTTT     |
| <i>Acs13</i>  | ACCACAGACATTAGCAGA     | AAGCACTCATTTTCATTCCT  |
| <i>Acc1</i>   | CCAGTCACACATACCCGAT    | TTCACACGACCCTAGGCT    |
| <i>Acc2</i>   | CACCCCTCCCAGAACCTC     | TCTCCACTACCCCAACATCCC |
| <i>Fas</i>    | CTACCTACCCGTGTGACCG    | CCACCAGAGACCGTTATGCC  |
| <i>Gpat1</i>  | ATCAGCAACATCATTCGGT    | AAAGCCCAAGAAGAACGAGA  |
| <i>Sreb1c</i> | TCTCAGTTGTTTCCTTTGCCTT | GATGCCGACCAGATTCCTT   |
| <i>Cpt1</i>   | AGATACACCAACGGGCTCA    | TGTCAAGAAAGGCACTCTCAC |
| <i>Ppara</i>  | AGGCCAAGTTGAAAGCAGAA   | GTCTTCTCTGCCATGCACAA  |