

Supplementary Table 1. Prevalence of fatty liver disease by hepatic steatosis index and simple nonalcoholic fatty liver disease score

Year	Overall		Male		Female	
	HSI >36	Simple NAFLD score ≥8	HSI >36	Simple NAFLD score ≥8	HSI >36	Simple NAFLD score ≥8
2009	46,476 (22.69)	63,589 (31.05)	28,151 (25.31)	39,204 (35.24)	18,325 (19.59)	24,385 (26.06)
2010	50,486 (22.23)	70,302 (30.95)	31,125 (25.4)	44,180 (36.06)	19,361 (18.50)	26,122 (24.96)
2011	51,431 (22.05)	73,629 (31.57)	31,445 (25.08)	45,735 (36.48)	19,986 (18.53)	27,894 (25.87)
2012	52,813 (21.90)	78,003 (32.35)	32,543 (25.14)	48,888 (37.76)	20,270 (18.15)	29,115 (26.07)
2013	54,894 (22.72)	79,961 (33.10)	33,720 (26.04)	49,679 (38.36)	21,174 (18.89)	30,282 (27.02)
2014	59,318 (22.83)	88,063 (33.90)	36,685 (26.43)	55,252 (39.81)	22,633 (18.70)	32,811 (27.11)
2015	63,361 (23.95)	94,866 (35.85)	39,112 (27.91)	59,624 (42.55)	24,249 (19.48)	35,242 (28.31)
2016	71,767 (24.91)	107,457 (37.30)	44,203 (29.10)	67,093 (44.17)	27,564 (20.24)	40,364 (29.63)
2017	74,166 (25.33)	113,422 (38.74)	45,892 (29.79)	71,002 (46.08)	28,274 (20.39)	42,420 (30.58)
<i>P</i> for trend	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

Values are presented as number (%).

HSI, hepatic steatosis index; NAFLD, nonalcoholic fatty liver disease.