

**Supplementary Table 3.** Clinical predictors for progression of rapid renal function decline in women according to menopause status

Variable	Premenopause ( <i>n</i> =176, age <55 yr)		Postmenopause ( <i>n</i> =236 age ≥55 yr)	
	OR (95% CI)	<i>P</i> value	OR (95% CI)	<i>P</i> value
Carotid plaque	0.46 (0.12–1.76)	0.256	2.12 (1.06–4.25)	0.034 <sup>a</sup>
Diabetes duration, yr	1.05 (0.96–1.15)	0.257	1.06 (1.01–1.12)	0.028 <sup>a</sup>
HbA1c, %	1.08 (0.80–1.45)	0.623	1.08 (0.88–1.33)	0.451
CVD	20.47 (1.70–245.83)	0.017 <sup>a</sup>	0.97 (0.37–2.52)	0.948
Hypertension	1.02 (0.37–2.85)	0.971	2.86 (1.35–6.07)	0.006 <sup>a</sup>
Use of ACEi/ARB	0.13 (0.01–1.25)	0.078	0.73 (0.34–1.60)	0.433

OR, odds ratio; CI, confidence interval; HbA1c, glycosylated hemoglobin; CVD, cardiovascular disease; ACEi, angiotensin-converting-enzyme inhibitor; ARB, angiotensin II-receptor blocker.

<sup>a</sup>Statistical significance.