

Supplementary Table 1. Baseline characteristics between controlled and uncontrolled diabetes

Characteristic	Controlled diabetes	Uncontrolled diabetes	P value
No. of patients	747	368	
Demographics			
Age, yr	55.7±7.5	55.8±7.6	0.887
Gender			
Men	627 (83.9)	295 (80.2)	0.117
Women	120 (16.1)	73 (19.8)	
Clinical characteristics or coexisting conditions			
Body mass index, kg/m ²	25.4±2.8	25.5±3.3	0.629
Obesity	405 (54.3)	194 (52.9)	0.653
Waist circumference, cm	88.7±7.6	89.1±8.4	0.424
Systolic blood pressure, mm Hg	122.8±13.3	124.4±14.3	0.059
Diastolic blood pressure, mm Hg	78.1±10.1	78.3±10.1	0.821
Hypertension	407 (54.5)	173 (47.0)	0.019
Hyperlipidemia	333 (44.6)	162 (44.0)	0.860
Current smoker	207 (27.7)	132 (35.9)	0.005
Previous stroke	13 (1.7)	4 (1.1)	0.402
Family history of CAD ^a	94 (12.6)	43 (11.7)	0.667
Fasting blood glucose, mg/dL	121.8±17.7	158.6±40.8	<0.001
HbA1c, %	6.2±0.5	8.1±1.2	<0.001
Total cholesterol, mg/dL	186.1±37.8	187.6±40.3	0.551
LDL-C, mg/dL	112.6±32.4	113.1±33.8	0.809
HDL-C, mg/dL	51.3±12.4	48.4±11.8	<0.001
Triglyceride, mg/dL	149.1±102.5	180.0±136.6	<0.001
Creatinine, mg/dL	0.9±0.1	0.9±0.2	0.011
Uric acid, mg/dL	5.7±1.3	5.3±1.3	<0.001
hs-CRP ≥2 mg/dL, %	9 (1.2)	7 (1.9)	0.357
Ejection fraction, %	63.4±4.0	63.6±4.4	0.518

Values are presented as mean ± standard deviation or number (%).

CAD, coronary artery disease; HbA1c, glycosylated hemoglobin; LDL-C, low density lipoprotein cholesterol; HDL-C, high density lipoprotein cholesterol; hs-CRP, high-sensitivity C-reactive protein.

^aCAD in a first-degree relative of any age.